



MIRROR

Reg no: B4873 Act 36 of 1947

Fertiliser Group 2

Active Ingredients:

Kaolin1 000 g/kg

Nett Mass / Volume

15 Kg

REGISTRATION HOLDER AND SUPPLIER DETAILS

Lutrasolve

2010/006368/07

51 Verdun Road, Kya Sands, 2196

011 792 8968

EMERGENCY CONTACT NUMBER

0861 METSON (638 766)

GHS (Pictogram)



WARNING / DANGER

BATCH NUMBER:

UN Number: Not regulated

Hazard Statements:

- Causes skin irritation.
- Causes serious eye irritation.
- Causes damage to organs through prolonged or repeated exposure.

Precautionary Statements:

- Poisonous when swallowed.
- Wear protective eye protection.
- Do not eat, drink, or smoke when mixing or using this product Do not inhale dust.
- Remove contaminated clothing immediately.
- IF ON SKIN: wash with plenty of soap and water.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and continue rinsing.
- In case of poisoning: CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM/HER.
- Store in a well-ventilated area. Keep container tightly closed.
- Destroy empty container by perforating, flattening and dispose of container in a safe way.

Growing with Metson/ Groei saam met ons



MIRROR

-
- Store away from food and feedstuffs.
 - Prevent contamination of food, feedstuffs, drinking water, eating utensils.
 - Keep out of reach of children.
 - The misuse of this product may cause serious damage to the crops. It should be used only under the direction of an agronomist. The manufacturer furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions that could not have been foreseen in terms of the label. Consult the supplier in the event of any uncertainty.

Directions for use: (use only as directed)

- Be advised to buffer the spray solution to a pH of 5.5.



MIRROR

Application Detail:

CROPS	SEASON LONG	LATE SEASON PROGRAM	COMMENTS
TREE CROPS			
1. Pome and Stone Fruit, Nut crops, Olives, Grape Vines, Small Fruit 2. Tropical fruit and Subtropical fruit (Avocados, Bananas, Mangos, Papaya, Pawpaw) 3. Nursery and Ornamental Plants 4. Citrus	1.2 kg / 100 L water beginning at petal fall. Follow-up every 14-21 days. Re-apply if rainfall and/or high temperatures occur.	1. Initial application 2.4kg/100L water. 2. Subsequently 1kg/100L water every 12 days. Cherries: Apply post- harvest only. Citrus: 1kg every 14-21 days with 4 applications.	Apply in water volumes according to the tree row volume. Prior to the first heat event. On hard wet foliage and grape vines add non-ionic surfactant at recommended label rates, the same at low rate of MIRROR and high water volume to the tree row volume. Use sprayers available depending on the size of the orchard and size of the trees
FIELD CROPS			
1. Peppers, Tomatoes, Potatoes, Onions, Cucurbits (Cucumber, Squash, Pumpkin, Melon and Watermelon) 2. Vegetable Seedlings 3. Pineapples 4. Cotton 5. Peanuts	1. 6.8kg/ha just prior to flowering 2. Re-apply at appropriate intervals according to the crop coverage required (generally every 14days) Sunburn and Heat Management 1. Initial application 25kg/ha Subsequent application 14kg/ha at the appropriate intervals according to crop coverage required		Apply the first 2 applications 10 days apart and prior to the first heat event. Increase the volume water used throughout the season based on the plant size. Re-apply at 10 days intervals as required to maintain an even coverage on the fruit and foliage. Maintain coverage up to 7 days before harvest. For transplant stock reduction apply to plant in trays. Cucurbits: apply only to smooth-skinned. Ensure that fruit is well covered for greatest sunburn reduction. DO NOT TREAT TRANSPLANT BY DRIPPING

Disclaimer / Conditions of Sale:

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors not controlled by the manufacturer. The registration holder furthermore does not accept responsibility for damage to crops, the environment, man or animal or for lack of performance of the remedy due to failure of the end user to follow label instructions or unforeseen events or conditions. Consult the supplier in the event of any uncertainty.



MIRROR



MIRROR

Reg no: B4873 Act 36 of 1947
FERTILISER GROUP 2

ACTIVE INGREDIENTS:

Kaolin

1 000 g/kg

DIRECTIONS FOR USE: Use only as directed

- Be advised to buffer the spray solution to a pH of 5.5.

APPLICATION DETAILS:

NETT MASS
VOLUME **15 kg**

REGISTRATION HOLDER: Lutrasolve
2010/006368/07
51 Verdun Road, Kya Sands, 2196
Tel: + 27 11 792 8968

EMERGENCY CONTACT: 0861 Metson (638 766)



WARNING / DANGER



Batch No:

Packed Date:

UN Number: Not Regulated



WARNING:
Keep away from food and feeds. Store in a cool dry place out of reach from children and/or uninformed persons and animals.

CROPS	SEASON LONG	LATE SEASON PROGRAM	COMMENTS
TREE CROPS			
1. Pome and Stone Fruit, Nut crops, Olives, Grape Vines, Small Fruit	1.2 kg / 100L water beginning at petal fall. Follow-up every 14-21 days. Re-apply if rainfall and/or high temperatures occur.	1. Initial application 2.4kg/100L water.	Apply in water volumes according to the tree row volume. Prior to the first heat event.
2. Tropical fruit and Subtropical fruit (Avocados, Bananas, Mangos, Papaya, Pawpaw)		2. Subsequently 1kg/100L water every 12 days.	On hard wet foliage and grape vines add non-ionic surfactant at recommended label rates, the same at low rate of MIRROR and high water volume to the tree row volume.
3. Nursery and Ornamental Plants		Cherries: Apply post-harvest only.	
4. Citrus		Citrus: 1kg every 14-21 days with 4 applications.	Use sprayers available depending on the size of the orchard and size of the trees